

OFIRMEV® (acetaminophen) for injection Reference Bibliography

This bibliography lists all identified manuscripts from PubMed of Randomized Controlled Clinical Trials and health economic and outcomes analyses (HEOR) studies, using intravenous acetaminophen (also known as paracetamol, especially outside the US)

Updated May 2019

US Basic Science

1. Raffa RB, Pawasauskas J, Pergolizzi JV Jr, Lu L, Chen Y, Wu S, Jarrett B, Fain R, Hill L, Devarakonda K. Pharmacokinetics of oral and intravenous paracetamol (acetaminophen) when co-administered with intravenous morphine in healthy adult subjects. *Clin Drug Investig.* 2018;38(3):259-268. [Link](#)
2. Singla NK, Parulan C, Samson R, Hutchinson J, Bushnell R, Beja EG, Ang R, Royal MA. Plasma and cerebrospinal fluid pharmacokinetic parameters after single-dose administration of intravenous, oral, or rectal acetaminophen. *Pain Pract.* 2012;12(7):523-32. [Link](#)

US Non-Surgical

3. Ankumah N-AE, Tsao M, Hutchinson M, Pedroza C, Mehta J, Sibai BM, Chauhan SP, Blackwell SC, Refuerzo JS. Intravenous acetaminophen versus morphine for analgesia in labor: a randomized trial. *Am J Perinatol.* 2017;34(1):38-43. [Link](#)
4. Barnaby DP, Chertoff AE, Restivo AJ, Campbell CM, Pearlman S, White D, Bijur PE, Gallagher EJ. Randomized controlled trial of intravenous acetaminophen versus intravenous hydromorphone for the treatment of acute pain in the emergency department. *Ann Emerg Med.* 2019; 73(2):133-140. [Link](#)
5. Candiotti KA, Bergese SD, Viscusi ER, Singla SK, Royal MA, Singla NK. Safety of multiple-dose intravenous acetaminophen in adult inpatients. *Pain Med.* 2010;11(12):1841-8. [Link](#)
6. Chang AK, Bijur PE, Ata A, Campbell C, Pearlman S, White D, Chertoff A, Restivo A, Gallagher EJ. Randomized clinical trial of IV acetaminophen as an analgesic adjunct for older adults with acute severe pain. *Acad Emerg Med.* 2019;26(4):402-409. [Link](#)
7. Kett DH, Breitmeyer JB, Ang R, Royal MA. A randomized study of the efficacy and safety of intravenous acetaminophen vs. intravenous placebo for the treatment of fever. *Clin Pharmacol Ther.* 2011;90(1):32-9. [Link](#)
8. Meyering SH, Stringer RW, Hysell MK. Randomized trial of adding parenteral acetaminophen to prochlorperazine and diphenhydramine to treat headache in the emergency department. *West J Emerg Med.* 2017;18(3):373-381. [Link](#)
9. Peacock WF, Breitmeyer JB, Pan C, Smith WB, Royal MA. A randomized study of the efficacy and safety of intravenous acetaminophen compared to oral acetaminophen for the treatment of fever. *Acad Emerg Med.* 2011;18(4):360-6. [Link](#)
10. Schell-Chaple HM, Liu KD, Matthay MA, Sessler DI, Puntillo KA. Effects of IV acetaminophen on core body temperature and hemodynamic responses in febrile critically ill adults: a randomized controlled trial. *Crit Care Med.* 2017;45(7):1199-1207 [Link](#)
11. Sin B, Wai M, Tatunchak T, Motov SM. The use of intravenous acetaminophen for acute pain in the emergency department. *Acad Emerg Med.* 2016;23(5):543-553. [Link](#)

US Surgical

12. Apfel CC, Souza K, Portillo J, Dalal P, Bergese SD. Patient satisfaction with intravenous acetaminophen: A pooled analysis of five randomized, placebo-controlled studies in the acute postoperative setting. *J Healthc Qual.* 2015;37(3):155-62. [Link](#)
13. Apfel CC, Turan A, Souza K, Pergolizzi J, Hornuss C. Intravenous acetaminophen reduces postoperative nausea and vomiting: A systematic review and meta-analysis. *Pain.* 2013;154(5):677-89. [Link](#)
14. Altenau B, Crisp CC, Devaiah CG, Lambers DS. Randomized controlled trial of intravenous acetaminophen for post-cesarean delivery pain control. *Am J Obstet Gynecol.* 217(3):362.e1-e362.e6. [Link](#)
15. Atencio I, Beushausen M, Kowalczyk JJ, Flores-Hidalgo A, Fino NF, Baur DA. Use of intravenous acetaminophen in postoperative pain management after partial and full bony impacted third molar extractions: a randomized double-blind controlled trial. *J Oral Maxillofac Surg.* 2018; 76(7):1414-1417. [Link](#)

16. Artime CA, Aijazi H, Zhang H, Syed T, Cai C, Gumbert SD, Ferrario L, C Normand K, Williams GW, Hagberg CA. Scheduled intravenous acetaminophen improves patient satisfaction with postcraniotomy pain management: a prospective, randomized, placebo-controlled, double-blind study. *J Neurosurg Anesthesiol.* 2018;30(3):231-236. [Link](#)
17. Aryaie AH, Lalezari S, Sergent WK, Puckett Y, Juergens C, Ratermann C, Ogg C. Decreased opioid consumption and enhance recovery with the addition of IV Acetaminophen in colorectal patients: a prospective, multi-institutional, randomized, double-blinded, placebo-controlled study (DOCIVA study). *Surg Endosc.* 2018;32(8):3432-3438. [Link](#)
18. Billings FT 4th, Petracek MR, Roberts LJ 2nd, Pretorius M. Perioperative intravenous acetaminophen attenuates lipid peroxidation in adults undergoing cardiopulmonary bypass: A randomized clinical trial. *PLoS One.* 2015;10(2):e0117625. [Link](#)
19. Blank JJ, Berger NG, Dux JP, Ali F, Ludwig KA, Peterson CY. The impact of intravenous acetaminophen on pain after abdominal surgery: a meta-analysis. *J Surg Res.* 2018;227:234-245. [Link](#)
20. Burbridge MA, Stone SA, Jaffe RA. Acetaminophen does not reduce postoperative opiate consumption in patients undergoing craniotomy for cerebral revascularization: a randomized control trial. *Cureus.* 2019;11(1):e3863 [Link](#)
21. Cooke FE, Samuels JD, Pomp A, Gadalla F, Wu X, Afaneh C, Dakin GF, Goldstein PA. A randomized, double-blind, placebo-controlled trial of intravenous acetaminophen on hospital length of stay in obese individuals undergoing sleeve gastrectomy. *Obes Surg.* 2018; 28(10):2998-3006. [Link](#)
22. Crisp CC, Khan M, Lambers DL, Westermann LB, Mazloomdoost DM, Yeung JJ, Kleeman SD, Pauls RN. The Effect of intravenous acetaminophen on postoperative pain and narcotic consumption after vaginal reconstructive surgery: a double-blind randomized placebo-controlled trial. *Female Pelvic Med Reconstr Surg.* 2017;23(2):80-85. [Link](#)
23. De Oliveira GSJ, Rodes ME, Bialek J, Kendall MC, McCarthy RJ. Single dose systemic acetaminophen to improve patient reported quality of recovery after ambulatory segmental mastectomy: A prospective, randomized, double-blinded, placebo controlled, clinical trial. *Breast J.* 2018;24(3):240-244. [Link](#)
24. De Oliveira GS Jr, Castro-Alves LJ, McCarthy RJ. Single-dose systemic acetaminophen to prevent postoperative pain: A meta-analysis of randomized controlled trials. *Clin J Pain.* 2015;31(1):86-93. [Link](#)
25. El Chaar M, Stoltzfus J, Claros L, Wasylik T. IV acetaminophen results in lower hospital costs and emergency room visits following bariatric surgery: A double-blind, prospective, randomized trial in a single accredited bariatric center. *J Gastrointest Surg.* 2016;20(4):715-24. [Link](#)
26. Fearon JA, Dimas V, Dittthakasem K, Herbert MA. A randomized controlled trial of oral versus intravenous administration of a nonnarcotic analgesia protocol following pediatric craniosynostosis corrections on nausea and vomiting rates. *J Craniofac Surg.* 2015;26(6):1951-3. [Link](#)
27. Greenberg S, Murphy GS, Avram MJ, Shear T, Benson J, Parikh KN, Patel A, Newmark R, Patel V, Bailes J, Szkol JW. Postoperative intravenous acetaminophen for craniotomy patients: a randomized controlled trial. *World Neurosurg.* 2018;109:e554-e562. [Link](#)
28. Gupta A, Abubaker H, Demas E, Ahrendtsen L. A randomized trial comparing the safety and efficacy of intravenous ibuprofen versus ibuprofen and acetaminophen in knee or hip arthroplasty. *Pain Physician.* 2016;19(6):349-356. [Link](#)
29. Hickman SR, Mathieson KM, Bradford LM, Garman CD, Gregg RW, Lukens DW. Randomized trial of oral versus intravenous acetaminophen for postoperative pain control. *Am J Health Syst Pharm.* 2018;75(6):367-375. [Link](#)
30. Jelacic S, Bollag L, Bowdle A, Rivat C, Cain KC, Richebe P. Intravenous acetaminophen as an adjunct analgesic in cardiac surgery reduces opioid consumption, but not opioid related adverse effects: a randomized controlled trial. *J Cardiothorac Vasc Anesth.* 2016;30(4):997-1004. [Link](#)
31. Lange M, Lee CW, Knisely T, Perla S, Barber K, Kia M. Efficacy of intravenous acetaminophen in length of stay and postoperative pain control in laparoscopic Roux-en-Y Gastric bypass surgery patients. *Bariatr Surg Pract Patient Care.* 2018;13(3):103-108. [Link](#)
32. Mamoun NF, Lin P, Zimmerman NM, Mascha EJ, Mick SL, Insler SR, Sessler DI, Duncan AE. Intravenous acetaminophen analgesia after cardiac surgery: A randomized, blinded, controlled superiority trial. *J Thorac Cardiovasc Surg.* 2016;152(3):881-889.e1. [Link](#).

33. Nour C, Ratsiu J, Singh N, Mason L, Ray A, Martin M, Hassanian M, Uhler J, Applegate RL. Analgesic effectiveness of acetaminophen for primary cleft palate repair in young children: A randomized placebo controlled trial. *Paediatr Anaesth*. 2014;24(6):574-81. [Link](#)
34. O'Neal JB, Freiberg AA, Yelle MD, Jiang Y, Zhang C, Gu Y, Kong X, Jian W, O'Neal WT, Wang J. Intravenous vs oral acetaminophen as an adjunct to multimodal analgesia after total knee arthroplasty: a prospective, randomized, double-blind clinical trial. *J Arthroplasty*. 2017;32(10):3029-3033. [Link](#)
35. Plunkett A, Haley C, McCoart A, Beltran T, Highland KB, Berry-Caban C, Lamberth S, Bartoszek M. A preliminary examination of the comparative efficacy of intravenous vs oral acetaminophen in the treatment of perioperative pain. *Pain Med*. 2017;18(12):2466-2473. [Link](#)
36. Politi JR, Davis RL 2nd, Matrka AK. Randomized prospective trial comparing the use of intravenous versus oral acetaminophen in total joint arthroplasty. *J Arthroplasty*. 2017;32(4):1125-1127. [Link](#)
37. Reagan KML, O'Sullivan DM, Gannon R, Steinberg AC. Decreasing postoperative narcotics in reconstructive pelvic surgery: a randomized controlled trial. *Am J Obstet Gynecol*. 2017;217(3):325.e1-e325.e10. [Link](#)
38. Rindos NB, Mansuria SM, Ecker AM, Stuparich MA, King CR. Intravenous acetaminophen versus saline in perioperative analgesia with laparoscopic hysterectomy. *Am J Obstet Gynecol*. 2019;220(4):373.e1-373.e8 [Link](#).
39. Rizkalla N, Zane NR, Prodell JL, Elci OU, Maxwell LG, DiLiberto MA, Zuppa AF. Use of intravenous acetaminophen in children for analgesia after spinal fusion surgery: a randomized clinical trial. *J Pediatr Pharmacol Ther*. 2018;23(5):395-404. [Link](#)
40. Roberts CA, Shah-Becker S, O'Connell Ferster A, Baker A, Stahl LE, Sedeek K, Carr MM. Randomized prospective evaluation of intraoperative intravenous acetaminophen in pediatric adenotonsillectomy. *Otolaryngol Head Neck Surg*. 2018;158(2):368-374. [Link](#)
41. Simpson SA, Zaccagni H, Bichell DP, et al. Acetaminophen attenuates lipid peroxidation in children undergoing cardiopulmonary bypass. *Pediatr Crit Care Med*. 2014;15(6):503-10. [Link](#)
42. Singla NK, Hale ME, Davis JC, et al. IV acetaminophen: Efficacy of a single dose for postoperative pain after hip arthroplasty: Subset data analysis of 2 unpublished randomized clinical trials. *Am J Ther*. 2015;22(1):2-10. [Link](#)
43. Sivakumar W, Jensen M, Martinez J, Tanana M, Duncan N, Hoesch R, Riva-Cambrin JK, Kilburg C, Ansari S, House PA. Intravenous acetaminophen for postoperative supratentorial craniotomy pain: a prospective, randomized, double-blinded, placebo-controlled trial. *J Neurosurg*. 2018;130(3):766-772. [Link](#)
44. Sola R Jr, Desai AA, Gonzalez KW, Doyle NM, Weaver KL, Poola AS, Fraser JD, St Peter SD, Millspaugh DL. Does intravenous acetaminophen improve postoperative pain control after laparoscopic appendectomy for perforated appendicitis? a prospective randomized trial. *Eur J Pediatr Surg*. 2019;29(2):159-165. [Link](#)
45. Strode MA, Sherman W, Mangieri CW, et al. Randomized trial of OFIRMEV versus placebo for pain management after laparoscopic sleeve gastrectomy. *Surg Obes Relat Dis*. 2016;12(4):772-777. [Link](#)
46. Suarez JC, Al-Mansoori AA, Kanwar S, Villa JM, McNamara CA, Patel PD. Effectiveness of novel adjuncts in pain management following total knee arthroplasty: a randomized clinical trial. *J Arthroplasty*. 2018;33(7S):S136-S141. [Link](#)
47. Subramaniam B, Shankar P, Shaefi S, Mueller A, O'Gara B, Banner-Goodspeed V, Gallagher J, Gasangwa D, Patxot M, Packiasabapathy S, Mathur P, Eikermann M, Talmor D, Marcantonio ER. Effect of intravenous acetaminophen vs placebo combined with propofol or dexmedetomidine on postoperative delirium among older patients following cardiac surgery: the DEXACET randomized clinical trial. *JAMA*. 2019;321(7):686-696. [Link](#)
48. Susheela AT, Packiasabapathy S, Gasangwa D-V, Patxot M, O'Neal J, Marcantonio E, Subramaniam B. The use of dexmedetomidine and intravenous acetaminophen for the prevention of postoperative delirium in cardiac surgery patients over 60 years of age: a pilot study. *F1000Res*. 2017;6:1842. [revised December 21, 2017]. [Link](#)
49. Takeda Y, Fukunishi S, Nishio S, Yoshiya S, Hashimoto K, Simura Y. Evaluating the effect of intravenous acetaminophen in multimodal analgesia after total hip arthroplasty: a randomized controlled trial. *J Arthroplasty*. 2019;[published February 25, 2019] [Link](#).
50. Thung AK, Elmaraghy CA, Barry N, Tumin D, Jatana KR, Rice J, Raman V, Bhalla T, Martin DP, Corridore M, Tobias JD. Double-blind randomized placebo-controlled trial of single-dose intravenous

- acetaminophen for pain associated with adenotonsillectomy in pediatric patients with sleep-disordered breathing. *J Pediatr Pharmacol Ther.* 2017;22(5):344-351. [Link](#)
51. Turan A, Karimi N, Zimmerman NM, Mick SL, Sessler DI, Mamoun N. Intravenous acetaminophen does not decrease persistent surgical pain after cardiac surgery. *J Cardiothorac Vasc Anesth.* 2017;31(6):2058-2064. [Link](#)
 52. Turner LC, Zyczynski HM, Shepherd JP. Intravenous acetaminophen before pelvic organ prolapse repair: a randomized controlled trial. *Obstet Gynecol.* 2019; 133(3):492-502 [Link](#)
 53. Towers CV, Shelton S, van Nes J, Gregory E, Liske E, Smalley A, Mobley E, Faircloth B, Fortner KB. Preoperative cesarean delivery intravenous acetaminophen treatment for postoperative pain control: a randomized double-blinded placebo control trial. *Am J Obstet Gynecol.* 2018;218(3):353.e1-353.e4. [Link](#)
 54. Tyler MA, Lam K, Ashoori F, Cai C, Kain JJ, Fakhri S, Citardi MJ, Cattano D, Luong A. Analgesic effects of intravenous acetaminophen vs placebo for endoscopic sinus surgery and postoperative pain: a randomized clinical trial. *JAMA Otolaryngol Head Neck Surg.* 2017;143(8):788-794 [Link](#)
 55. Viswanath A, Oreadi D, Finkelman M, Klein G, Papageorge M. Does pre-emptive administration of intravenous ibuprofen (Caldolor) or intravenous acetaminophen (Ofirmev) reduce postoperative pain and subsequent narcotic consumption after third molar surgery? *J Oral Maxillofac Surg.* 2019; 77(2):262-270. [Link](#)
 56. Wang VC, Preston MA, Kibel AS, Xu X, Gosnell J, Yong RJ, Urman RD. A prospective, randomized, double-blind, placebo-controlled trial to evaluate intravenous acetaminophen versus placebo in patients undergoing robotic-assisted laparoscopic prostatectomy. *J Pain Palliat Care Pharmacother.* 2019;[published online: January 15, 2019]; [Link](#).
 57. Westrich GH, Birch GA, Muskat AR, Padgett DE, Goytizolo EA, Bostrom MP, Mayman DJ, Lin Y, YaDeau JT. Intravenous vs oral acetaminophen as a component of multimodal analgesia after total hip arthroplasty: a randomized, blinded trial. *J Arthroplasty.* 2019;[published online: March 6, 2019] [Link](#).
 58. Wilson SH, Wolf BJ, Robinson SM, Nelson C, Hebbard L. Intravenous vs oral acetaminophen for analgesia after cesarean delivery: a randomized trial. *Pain Med.* 2018; [published online: December 17, 2018]. [Link](#)
 59. Winger SJ, Miller H, Minkowitz HS, Royal MA, Ang RY, Breitmeyer JB, Singla NK. A randomized, double-blind, placebo-controlled, multicenter, repeat-dose study of two intravenous acetaminophen dosing regimens for the treatment of pain after abdominal laparoscopic surgery. *Clin Ther.* 2010;32(14):2348-69. [Link](#)
 60. Ziemann-Gimmel P, Goldfarb AA, Koppman J, Marema RT. Opioid-free total intravenous anaesthesia reduces postoperative nausea and vomiting in bariatric surgery beyond triple prophylaxis. *Br J Anaesth.* 2014;112(5):906-11. [Link](#)

Outside US – Basic Sciences

61. Chiam E, Weinberg L, Bailey M, McNicol L, Bellomo R. The haemodynamic effects of intravenous paracetamol (acetaminophen) in healthy volunteers: A double-blinded, randomized, triple crossover trial. *Br J Clin Pharmacol.* 2016;81(4):605-12. [Link](#)
62. Flint RB, Roofthoof DW, van Rongen A, van Lingen RA, van den Anker JN, van Dijk M, Allegaert K, Tibboel D, Knibbe CAJ, Simons SHP. Exposure to acetaminophen and all its metabolites upon 10, 15, and 20 mg/kg intravenous acetaminophen in very-preterm infants. *Pediatr Res.* 2017;82(4):678-684. [Link](#)
63. Pickering G, Moustafa F, Desbrandes S, Cardot JM, Roux D, Dubray C. Paracetamol and opioid pathways: A pilot randomized clinical trial. *Fundam Clin Pharmacol.* 2013;27(3):339-45. [Link](#)

Outside US – Non-Surgical

64. Abd-El-Maeboud KH, Elboholy AE, Mohammed WE, Elgamel HM, Ali WA. Intravenous infusion of paracetamol for intrapartum analgesia. *J Obstet Gynaecol Res.* 2014;40(11):2152-7. [Link](#)
65. Abd El-Mashad E-R, El-Mahdy H, El Amrousy D, Elgendy M. Comparative study of the efficacy and safety of paracetamol, ibuprofen, and indomethacin in closure of patent ductus arteriosus in preterm neonates. *Eur J Pediatr.* 2017;176(2):233-240. [Link](#)
66. Abdollahi MH, Mojibian M, Pishgahi A, Mallah F, Dareshiri S, Mohammadi S, Naghavi-Behzad M. Intravenous paracetamol versus intramuscular pethidine in relief of labour pain in primigravid women. *Niger Med J.* 2014;55(1):54-7. [Link](#)

67. Ahmadi A, Amri P, Shokri J, Hajian K. Comparison of the analgesic effect of intravenous paracetamol/midazolam and fentanyl in preparation of patients for colonoscopy: A double blind randomized clinical trial. *Caspian J Intern Med.* 2015;6(2):87-92. [Link](#)
68. Akinci N, Bakan N, Karaoren G, Tomruk SG, Sokmen HM, Yanli Y, Akcay ME. Comparison of Clinical Effects of Dexketoprofen and Paracetamol Used for Analgesia in Endoscopic Retrograde Cholangiopancreatography. *Turk J Anaesthesiol Reanim.* 2016;44(1):13-20. [Link](#).
69. Aksel G, Guler S, Dogan NO, Corbacioglu SK. A randomized trial comparing intravenous paracetamol, topical lidocaine, and ice application for treatment of pain associated with scorpion stings. *Hum Exp Toxicol.* 2015;34(6):662-7. [Link](#)
70. Al B, Sunar MM, Zengin S, Sabak M, Bogan M, Can B, Kul S, Murat Oktay M, Eren SH. Comparison of intravenous dexketoprofen trometamol, fentanyl, and paracetamol in the treatment of patients admitted to the emergency department for renal colic: A randomized controlled trial. *Am J Emerg Med.* 2018;36(4):571-576. [Link](#)
71. Azizkhani R, Pourafzali SM, Baloochestani E, Masoumi B. Comparing the analgesic effect of intravenous acetaminophen and morphine on patients with renal colic pain referring to the emergency department: A randomized controlled trial. *J Res Med Sci.* 2013;18(9):772-6. [Link](#)
72. Bektas F, Eken C, Karadeniz O, Goksu E, Cubuk M, Cete Y. Intravenous paracetamol or morphine for the treatment of renal colic: A randomized, placebo-controlled trial. *Ann Emerg Med.* 2009;54(4):568-74. [Link](#)
73. Borazan H, Erdem TB, Kececioglu M, Otelcioglu S. Prevention of pain on injection of propofol: A comparison of lidocaine with different doses of paracetamol. *Eur J Anaesthesiol.* 2010;27(3):253-7. [Link](#)
74. Canbay O, Celebi N, Arun O, Karagoz AH, Saricaoglu F, Ozgen S. Efficacy of intravenous acetaminophen and lidocaine on propofol injection pain. *Br J Anaesth.* 2008;100(1):95-8. [Link](#)
75. Cenker E, Serinken M, Uyanik E. Intravenous paracetamol vs ibuprofen in renal colic: a randomised, double-blind, controlled clinical trial. *Urolithiasis.* 2018; 46(4):369-373. [Link](#)
76. Craig M, Jeavons R, Probert J, Bengler J. Randomised comparison of intravenous paracetamol and intravenous morphine for acute traumatic limb pain in the emergency department. *Emerg Med J.* 2012;29(1):37-9. [Link](#)
77. Cuvillon P, Zoric L, Demattei C, Alonso S, Casano F, L'hermite J, Ripart J, Lefrant J-Y, Muller L. Opioid-sparing effect of nefopam in combination with paracetamol after major abdominal surgery: a randomized double-blind study. *Minerva Anesthesiol.* 2017;83(9):914-920. [Link](#)
78. Demirozogul E, Yilmaz A, Ozen M, Turkcuer I, Seyit M, Arikan C. Intravenous dexketoprofen versus paracetamol in non-traumatic musculoskeletal pain in the emergency department: a randomized clinical trial. *Am J Emerg Med.* 2019;[published online: February 4, 2019]. [Link](#)
79. Duhamel JF, Le Gall E, Dalphin ML, Payen-Champenois C. Antipyretic efficacy and safety of a single intravenous administration of 15 mg/kg paracetamol versus 30 mg/kg propacetamol in children with acute fever due to infection. *Int J Clin Pharmacol Ther.* 2007;45(4):221-9. [Link](#)
80. Eken C, Serinken M, Elicabuk H, Uyanik E, Erdal M. Intravenous paracetamol versus dexketoprofen versus morphine in acute mechanical low back pain in the emergency department: A randomised double-blind controlled trial. *Emerg Med J.* 2014;31(3):177-81. [Link](#)
81. Elbohuty AE, Abd-Elrazek H, Abd-El-Gawad M, Salama F, El-Shorbagy M, Abd-El-Maeboud KH. Intravenous infusion of paracetamol versus intravenous pethidine as an intrapartum analgesic in the first stage of labor. *Int J Gynaecol Obstet.* 2012;118(1):7-10. [Link](#)
82. Esmailian M, Moshiri R, Zamani M. Comparison of the analgesic effect of intravenous acetaminophen and morphine sulfate in rib fracture; a randomized double-blind clinical trial. *Emerg (Tehran).* 2015;3(3):99-102. [Link](#)
83. Evron S, Ezri T, Protianov M, et al. The effects of remifentanyl or acetaminophen with epidural ropivacaine on body temperature during labor. *J Anesth.* 2008;22(2):105-11. [Link](#)
84. Faridaalae G, Rahmani SH, Mehryar H, et al. Comparison of intravenous metoclopramide and acetaminophen in primary headaches: A randomized controlled trial. *Emerg (Tehran).* 2015;3(2):70-4. [Link](#)
85. Farnia MR, Babaei R, Shirani F, et al. Analgesic effect of paracetamol combined with low-dose morphine versus morphine alone on patients with biliary colic: A double blind, randomized controlled trial. *World J Emerg Med.* 2016;7(1):25-9. [Link](#)

86. Filitz J, Ihmsen H, Gunther W, et al. Supra-additive effects of tramadol and acetaminophen in a human pain model. *Pain*. 2008;136(3):262-70. [Link](#)
87. Furyk J, Levas D, Close B, Laspina K, Fitzpatrick M, Robinson K, Vangaveti VN, Ray R. Intravenous versus oral paracetamol for acute pain in adults in the emergency department setting: a prospective, double-blind, double-dummy, randomised controlled trial. *Emerg Med J*. 2018;35(3):179-184. [Link](#)
88. Ghaderian M, Armanian AM, Sabri MR, Montaseri M. Low-dose intravenous acetaminophen versus oral ibuprofen for the closure of patent ductus arteriosus in premature neonates. *J Res Med Sci*. 2019;24:13. [Link](#).
89. Grissa MH, Claessens YE, Boudia W, et al. Paracetamol vs piroxicam to relieve pain in renal colic. results of a randomized controlled trial. *Am J Emerg Med*. 2011;29(2):203-6. [Link](#)
90. Gulen B, Dur A, Serinken M, Karcioğlu O, Sonmez E. Pain treatment in patients with acute pancreatitis: A randomized controlled trial. *Turk J Gastroenterol*. 2016;27(2):192-6. [Link](#)
91. Gupta K, Mitra S, Kazal S, Saroa R, Ahuja V, Goel P. I.V. paracetamol as an adjunct to patient-controlled epidural analgesia with levobupivacaine and fentanyl in labour: a randomized controlled study. *Br J Anaesth*. 2016;117(5):617-622. [Link](#)
92. Harkin P, Harma A, Aikio O, Valkama M, Leskinen M, Saarela T, Hallman M. Paracetamol accelerates closure of the ductus arteriosus after premature birth: a randomized trial. *J Pediatr*. 2016;177:72-77.e2. [Link](#).
93. Hochwald O, Mainzer G, Borenstein-Levin L, Jubran H, Dinur G, Zucker M, Mor M, Khoury A, Kugelman A. adding paracetamol to Ibuprofen for the treatment of patent ductus arteriosus in preterm infants: a double-blind, randomized, placebo-controlled pilot study. *Am J Perinatol*. 2018;35(13):1319-1325. [Link](#)
94. Honarmand H, Abdollahi M, Ahmadi A, et al. Randomized trial of the effect of intravenous paracetamol on inflammatory biomarkers and outcome in febrile critically ill adults. *Daru*. 2012;20(1):12. [Link](#)
95. Huang X, Wang F, Wang K. Paracetamol versus ibuprofen for the treatment of patent ductus arteriosus in preterm neonates: a meta-analysis of randomized controlled trials. *J Matern Fetal Neonatal Med*. 2018;31(16):2216-2222. [Link](#)
96. Imani F, Faiz HR, Sedaghat M, Hajiashrafi M. Effects of adding ketamine to fentanyl plus acetaminophen on postoperative pain by patient controlled analgesia in abdominal surgery. *Anesth Pain Med*. 2014;4(1):e12162. [Link](#)
97. Ing Lorenzini K, Besson M, Daali Y, Salomon D, Dayer P, Desmeules J. A randomized, controlled trial validates a peripheral supra-additive antihyperalgesic effect of a paracetamol-ketorolac combination. *Basic Clin Pharmacol Toxicol*. 2011;109(5):357-64. [Link](#)
98. Jalili M, Mozaffarpour Noori A, Sedaghat M, Safaie A. Efficacy of intravenous paracetamol versus intravenous morphine in acute limb trauma. *Trauma Mon*. 2016;21(1):e19649. [Link](#)
99. Jeon Y, Baek SU, Park SS, Kim SO, Baek WY, Yeo JS. Effect of pretreatment with acetaminophen on withdrawal movements associated with injection of rocuronium: A prospective, randomized, double-blind, placebo controlled study. *Korean J Anesthesiol*. 2010;59(1):13-6. [Link](#)
100. Juujarvi S, Kallankari H, Patsi P, Leskinen M, Saarela T, Hallman M, Aikio O. Follow-up study of the early, randomised paracetamol trial to preterm infants, found no adverse reactions at the two-years corrected age. *Acta Paediatr*. 2019;108(3):452-458. [Link](#)
101. Kaur Makkar J, Jain K, Bhatia N, Jain V, Mal Mithrawal S. Comparison of analgesic efficacy of paracetamol and tramadol for pain relief in active labor. *J Clin Anesth*. 2015;27(2):159-63. [Link](#)
102. Kaynar M, Koyuncu F, Buldu I, et al. Comparison of the efficacy of diclofenac, acupuncture, and acetaminophen in the treatment of renal colic. *Am J Emerg Med*. 2015;33(6):749-53. [Link](#)
103. Kelly SJ, Moran JL, Williams PJ, Burns K, Rowland A, Miners JO, Peake SL. Haemodynamic effects of parenteral vs. enteral paracetamol in critically ill patients: a randomised controlled trial. *Anaesthesia*. 2016;71(10):1153-1162. [Link](#)
104. Kouček M, Mansouri B, Mokhtari M, Goharani R, Miri MM, Sistanizad M. A comparative study of intravenous paracetamol and fentanyl for pain management in ICU. *Iran J Pharm Res*. 2013;12(1):193-8. [Link](#)
105. Lallar M, Anam HU, Nandal R, Singh SP, Katyal S. Intravenous paracetamol infusion versus intramuscular tramadol as an intrapartum labor analgesic. *J Obstet Gynaecol India*. 2015;65(1):17-22. [Link](#)

106. Leinisch E, Evers S, Kaempfe N, Sostak P, Jürgens T, Straube A, May A. Evaluation of the efficacy of intravenous acetaminophen in the treatment of acute migraine attacks: A double-blind, placebo-controlled parallel group multicenter study. *Pain*. 2005;117(3):396-400. [Link](#)
107. Manne VSSK, Gondi SR. Comparative Study of the Effect of intravenous paracetamol and tramadol in relieving of postoperative pain after general anesthesia in nephrectomy patients. *Anesth Essays Res*. 2017;11(1):117-120. [Link](#)
108. Masoumi K, Forouzan A, Asgari Darian A, Feli M, Barzegari H, Khavanin A. Comparison of clinical efficacy of intravenous acetaminophen with intravenous morphine in acute renal colic: a randomized, double-blind, controlled trial. *Emerg Med Int*. 2014;2014:571326. [Link](#)
109. Medina-Vera AJ, Novoa LM. Reduced anaesthetic requirements and postoperative analgesics in patients undergoing laparoscopic cholecystectomy: premedication with intravenous paracetamol versus ketorolac, a double blind and randomised clinical trial. *Rev Esp Anesthesiol Reanim*. 2017;64(2):64-70. [Link](#)
110. Momeni M, Vahidi E, Badrizadeh M, Naderpour Z, Saeedi M. Oral diclofenac potassium versus intravenous acetaminophen in acute, isolated, closed-limb trauma. *Adv Emerg Nurs J*. 2019;41(1):48-55. [Link](#)
111. Munsterhjelm E, Niemi TT, Ylikorkala O, Neuvonen PJ, Rosenberg PH. Influence on platelet aggregation of I.V. parecoxib and acetaminophen in healthy volunteers. *Br J Anaesth*. 2006;97(2):226-31. [Link](#)
112. Paramba FC, Naushad VA, Purayil N, Mohammed OH, Chandra P. Randomized controlled study of the antipyretic efficacy of oral paracetamol, intravenous paracetamol, and intramuscular diclofenac in patients presenting with fever to the emergency department. *Ther Clin Risk Manag*. 2013;9:371-6. [Link](#)
113. Parker SL, Saxena M, Gowardman J, Lipman J, Myburgh J, Roberts JA. Population pharmacokinetics of intravenous paracetamol in critically ill patients with traumatic brain injury. *J Crit Care*. 2018;47:15-20. [Link](#)
114. Pathan SA, Mitra B, Straney LD, et al. Delivering safe and effective analgesia for management of renal colic in the emergency department: A double-blind, multigroup, randomised controlled trial. *Lancet*. 2016;387(10032):1999-2007. [Link](#)
115. Pickering G, Moustafa F, Macian N, Schmidt J, Pereira B, Dubray C. A new transmucous-buccal formulation of acetaminophen for acute traumatic pain: A non-inferiority, randomized, double-blind, clinical trial. *Pain Physician*. 2015;18(3):249-57. [Link](#)
116. Roy S, Simalti AK. Comparison of antipyretic efficacy of intravenous (IV) acetaminophen versus oral (PO) acetaminophen in the management of fever in children. *Indian J Pediatr*. 2018;85(1):1-4. [Link](#)
117. Saxena MK, Taylor C, Billot L, Bompont S, Gowardman J, Roberts JA, Lipman J, Myburgh JI. The effect of paracetamol on core body temperature in acute traumatic brain injury: A randomised, controlled clinical trial. *PLoS One*. 2015;10(12):e0144740. [Link](#)
118. Scharbert G, Gebhardt K, Sow Z, Duris M, Deusch E, Kozek-Langenecker S. Point-of-care platelet function tests: Detection of platelet inhibition induced by nonopioid analgesic drugs. *Blood Coagul Fibrinolysis*. 2007;18(8):775-80. [Link](#)
119. Serinken M, Eken C, Gungor F, Emet M, Al B. Comparison of intravenous morphine vs paracetamol in sciatica: A randomized placebo controlled trial. *Acad Emerg Med*. 2016;23(6):674-678. [Link](#)
120. Serinken M, Eken C, Turkcuer I, Elicabuk H, Uyanik E, Schultz CH. Intravenous paracetamol versus morphine for renal colic in the emergency department: A randomised double-blind controlled trial. *Emerg Med J*. 2012;29(11):902-5. [Link](#)
121. Serinken M, Eken C, Karcioğlu O. IV dexametopfen vs. IV paracetamol in patients presented with dysmenorrhea to emergency department: a randomized controlled trial. *Balkan Med J*. 2018; 35(4):301-305. [Link](#)
122. Shams Vahdati S, Morteza Baghi HR, Ghobadi J, Rajaei Ghafouri R, Habibollahi P. Comparison of paracetamol (apotel(R)) and morphine in reducing post pure head trauma headache. *Anesth Pain Med*. 2014;4(3):e14903. [Link](#)
123. Tasmacioglu B, Aydinli I, Keskinbora K, Pekel AF, Salihoglu T, Sonsuz A. Effect of intravenous administration of paracetamol on morphine consumption in cancer pain control. *Support Care Cancer*. 2009;17(12):1475-81. [Link](#)
124. Tsaganos T, Tseti IK, Tziolos N, Soumelas G-S, Koupetori M, Pырpasopoulou A, Akinosoglou K, Gogos C, Tsokos N, Karagiannis A, Sympardi S, Giamarellos-Bourboulis EJ. Randomized controlled

- multicenter clinical trial of the antipyretic effect of intravenous paracetamol in patients admitted to hospital with infection. *Br J Clin Pharmacol.* 2017;83(4):742-750. [Link](#)
125. Turkcuer I, Serinken M, Eken C, Yilmaz A, Akdag Ö, Uyan E, Kiray C, Elicabuk H. Intravenous paracetamol versus dexketoprofen in acute migraine attack in the emergency department: A randomised clinical trial. *Emerg Med J.* 2014;31(3):182-5. [Link](#)
126. Yilmaz A, Sabirli R, Ozen M, Turkcuer I, Erdur B, Arikan C, Demirozogul E, Sarohan A, Seyit M, Ok N. Intravenous paracetamol versus dexketoprofen in acute musculoskeletal trauma in the emergency department: A randomised clinical trial. *Am J Emerg Med.* 2019; 37(5):902-908. [Link](#)
127. Young P, Saxena M, Bellomo R, Freebairn R, Hammond N, van Haren F, Holliday M, Henderson S, Mackle D, McArthur C, McGuinness S, Myburgh J, Weatherall M, Webb , Beasley R; HEAT Investigators; Australian and New Zealand Intensive Care Society Clinical Trials Group. Acetaminophen for fever in critically ill patients with suspected infection. *N Engl J Med.* 2015;373(23):2215-24. [Link](#)
128. Zare MA, Ghalyaie AH, Fathi M, Farsi D, Abbasi S, Hafezimoghadam P. Oral oxycodone plus intravenous acetaminophen versus intravenous morphine sulfate in acute bone fracture pain control: A double-blind placebo-controlled randomized clinical trial. *Eur J Orthop Surg Traumatol.* 2014;24(7):1305-9. [Link](#)

Outside US – Surgical

129. Abdulla S, Eckhardt R, Netter U, Abdulla W. A randomized, double-blind, controlled trial on non-opioid analgesics and opioid consumption for postoperative pain relief after laparoscopic cholecystectomy. *Acta Anaesthesiol Belg.* 2012;63(1):43-50. [Link](#)
130. Abdulla S, Eckhardt R, Netter U, Abdulla W. Efficacy of three IV non-opioid-analgesics on opioid consumption for postoperative pain relief after total thyroidectomy: A randomised, double-blind trial. *Middle East J Anaesthesiol.* 2012;21(4):543-52. [Link](#)
131. Abdulla S, Eckhardt R, Netter U, Abdulla W. Randomized, double-blind, placebo-controlled study to assess the efficacy of nonopioid analgesics on pain following arthroscopic knee surgery. *Pain Res Treat.* 2012;2012:305821. [Link](#)
132. Acmaz G, Aksoy H, Ozoglu N, Aksoy U, Albayrak E. Effect of paracetamol, dexketoprofen trometamol, lidocaine spray, and paracervical block application for pain relief during suction termination of first-trimester pregnancy. *Biomed Res Int.* 2013;2013:869275. [Link](#)
133. Akcali GE, Iskender A, Demiraran Y, et al. Randomized comparison of efficacy of paracetamol, lornoxicam, and tramadol representing three different groups of analgesics for pain control in extracorporeal shockwave lithotripsy. *J Endourol.* 2010;24(4):615-20. [Link](#)
134. Akil A, Api O, Bektas Y, Yilmaz AO, Yalti S, Unal O. Paracetamol vs dexketoprofen for perineal pain relief after episiotomy or perineal tear. *J Obstet Gynaecol.* 2014;34(1):25-8. [Link](#)
135. Aksoy M, Ince I, Ahiskalioglu A, Keles S, Doymus O. Effect of intravenous preoperative versus postoperative paracetamol on postoperative nausea and vomiting in patients undergoing strabismus surgery: A prospective randomized study. *Agri.* 2018;30(1):1-7. [Link](#)
136. Alhashemi JA, Daghistani MF. Effect of intraoperative intravenous acetaminophen vs. intramuscular meperidine on pain and discharge time after paediatric dental restoration. *Eur J Anaesthesiol.* 2007;24(2):128-33. [Link](#)
137. Alhashemi JA, Alotaibi QA, Mashaat MS, Kaid TM, Mujallid RH, Kaki AM. Intravenous acetaminophen vs oral ibuprofen in combination with morphine PCIA after cesarean delivery. *Can J Anaesth.* 2006;53(12):1200-6. [Link](#)
138. Alhashemi JA, Daghistani MF. Effects of intraoperative I.V. acetaminophen vs I.M. meperidine on post-tonsillectomy pain in children. *Br J Anaesth.* 2006;96(6):790-5. [Link](#)
139. Ali MA, Shamim F, Chughtai S. Comparison between intravenous paracetamol and fentanyl for intraoperative and postoperative pain relief in dilatation and evacuation: Prospective, randomized interventional trial. *J Anaesthesiol Clin Pharmacol.* 2015;31(1):54-8. [Link](#)
140. Api O, Unal O, Ugurel V, Emeksiz MB, Turan C. Analgesic efficacy of intravenous paracetamol for outpatient fractional curettage: A randomised, controlled trial. *Int J Clin Pract.* 2009;63(1):105-11. [Link](#)
141. Arici S, Gurbet A, Turker G, Yavascaoglu B, Sahin S. Preemptive analgesic effects of intravenous paracetamol in total abdominal hysterectomy. *Agri.* 2009;21(2):54-61. [Link](#)

142. Arslan M, Celep B, Cicek R, Kalender HU, Yilmaz H. Comparing the efficacy of preemptive intravenous paracetamol on the reducing effect of opioid usage in cholecystectomy. *J Res Med Sci.* 2013;18(3):172-7. [Link](#)
143. Arslan M, Cicek R, Celep B, Yilmaz H, Ustun Kalender H. [Comparison of the analgesic effects of intravenous paracetamol and lornoxicam in postoperative pain following thyroidectomies]. *Agri.* 2011;23(4):160-6. [Link](#)
144. Atef A, Fawaz AA. Intravenous paracetamol is highly effective in pain treatment after tonsillectomy in adults. *Eur Arch Otorhinolaryngol.* 2008;265(3):351-5. [Link](#)
145. Ayatollahi V, Faghihi S, Behdad S, Heiranizadeh N, Baghianimoghadam B. Effect of preoperative administration of intravenous paracetamol during cesarean surgery on hemodynamic variables relative to intubation, postoperative pain and neonatal apgar. *Acta Clin Croat.* 2014;53(3):272-8. [Link](#)
146. Bameshki A, Peivandi Yazdi A, Sheybani S, Rezaei Boroujerdi H, Taghavi Gilani M. The assessment of addition of either intravenous paracetamol or diclofenac suppositories to patient-controlled morphine analgesia for postgastrectomy pain control. *Anesth Pain Med.* 2015;5(5):e29688. [Link](#)
147. Baygin O, Tuzuner T, Isik B, Kusgoz A, Tanriver M. Comparison of pre-emptive ibuprofen, paracetamol, and placebo administration in reducing post-operative pain in primary tooth extraction. *Int J Paediatr Dent.* 2011;21(4):306-13. [Link](#)
148. Bilir S, Yurtlu BS, Hanci V, Okyay RD, Erdogan Kayhan G, Ayoglu HP, Ozkocak Turan I. Effects of peroperative intravenous paracetamol and lornoxicam for lumbar disc surgery on postoperative pain and opioid consumption: A randomized, prospective, placebo-controlled study. *Agri.* 2016;28(2):98-105. [Link](#).
149. Bonnal A, Dehon A, Nagot N, Macioce V, Nogue E, Morau E. Patient-controlled oral analgesia versus nurse-controlled parenteral analgesia after caesarean section: a randomised controlled trial. *Anaesthesia.* 2016;71(5):535-543. [Link](#)
150. Borisov DB, Levin AV, Uvarov DN, Kapanadze LG, Nedashkovskii EV. [Balanced postoperative analgesia in abdominal surgery: Efficiency of the combined use of epidural block and non-opioid analgesics]. *Anesteziol Reanimatol.* 2009;-2(2):35-7. [Link](#)
151. Borisov DB, Levin AV, Vyl'iuov IV, Sokolov AV, Nedashkovskii EV. [Efficiency of preemptive intravenous paracetamol analgesia in abdominal surgery]. *Anesteziol Reanimatol.* 2007;-5(5):38-40. [Link](#)
152. Brett CN, Barnett SG, Pearson J. Postoperative plasma paracetamol levels following oral or intravenous paracetamol administration: A double-blind randomised controlled trial. *Anaesth Intensive Care.* 2012;40(1):166-71. [Link](#)
153. Brodner G, Gogarten W, Van Aken H, et al. Efficacy of intravenous paracetamol compared to dipyron and parecoxib for postoperative pain management after minor-to-intermediate surgery: A randomised, double-blind trial. *Eur J Anaesthesiol.* 2011;28(2):125-32. [Link](#)
154. Cakan T, Inan N, Culhaoglu S, Bakkal K, Basar H. Intravenous paracetamol improves the quality of postoperative analgesia but does not decrease narcotic requirements. *J Neurosurg Anesthesiol.* 2008;20(3):169-73. [Link](#)
155. Caliskan E, Sener M, Kocum A, Ozyilkan NB, Ezer SS, Aribogan A. The efficacy of intravenous paracetamol versus dipyron for postoperative analgesia after day-case lower abdominal surgery in children with spinal anesthesia: A prospective randomized double-blind placebo-controlled study. *BMC Anesthesiol.* 2013;13(1):34. [Link](#)
156. Caliskan E, Sener M, Kipri M, Yilmaz I, Aribogan A. Comparison of the effects of intravenous dexketoprofen trometamol versus paracetamol on postoperative analgesia in patients undergoing septoplasty: A randomised double-blind clinical trial. *Pak J Med Sci.* 2018;34(3):546-552. [Link](#)
157. Capici F, Ingelmo PM, Davidson A, et al. Randomized controlled trial of duration of analgesia following intravenous or rectal acetaminophen after adenotonsillectomy in children. *Br J Anaesth.* 2008;100(2):251-5. [Link](#)
158. Cattabriga I, Pacini D, Lamazza G, et al. Intravenous paracetamol as adjunctive treatment for postoperative pain after cardiac surgery: A double blind randomized controlled trial. *Eur J Cardiothorac Surg.* 2007;32(3):527-31. [Link](#)
159. Ceelie I, de Wildt SN, van Dijk M, et al. Effect of intravenous paracetamol on postoperative morphine requirements in neonates and infants undergoing major noncardiac surgery: A randomized controlled trial. *JAMA.* 2013;309(2):149-54. [Link](#)

160. Caliskan E, Sener M, Kipri M,, Yilmaz I, Aribogan A. Comparison of the effects of intravenous Dexketoprofen Trometamol versus Paracetamol on postoperative analgesia in patients undergoing Septoplasty: A randomised double-blind clinical trial. *Pak J Med Sci.* 2018;34(3):546-552. [Link](#)
161. Celik EC, Kara D, Koc E, Yayik AM. The comparison of single-dose preemptive intravenous ibuprofen and paracetamol on postoperative pain scores and opioid consumption after open septorhinoplasty: a randomized controlled study. *Eur Arch Otorhinolaryngol.* 2018; 275(9):2259-2263. [Link](#)
162. Ceyhan D, Bilir A, Gulec MS. Evaluation of the analgesic efficacy of dexketoprofen added to paracetamol. *Turk anestezi reanim.* 2016;44(6):312-316. [Link](#)
163. Chiam E, Bellomo R, Churilov L, Weinberg L. The hemodynamic effects of intravenous paracetamol (acetaminophen) vs normal saline in cardiac surgery patients: A single center placebo controlled randomized study. *PLoS One.* 2018;13(4):e0195931. [Link](#).
164. Choudhuri AH, Uppal R. A comparison between intravenous paracetamol plus fentanyl and intravenous fentanyl alone for postoperative analgesia during laparoscopic cholecystectomy. *Anesth Essays Res.* 2011;5(2):196-200. [Link](#)
165. Ciftci B, Ekinci M, Celik EC, Kaciroglu A, Karakaya MA, Demiraran Y, Ozdenkaya Y. Comparison of intravenous ibuprofen and paracetamol for postoperative pain management after laparoscopic sleeve gastrectomy. a randomized controlled study. *Obes Surg.* 2019;29(3):765-770. [Link](#).
166. Cok OY, Eker HE, Pelit A, et al. The effect of paracetamol on postoperative nausea and vomiting during the first 24 h after strabismus surgery: A prospective, randomised, double-blind study. *Eur J Anaesthesiol.* 2011;28(12):836-41. [Link](#)
167. Cornesse D, Senard M, Hans GA, et al. Comparison between two intraoperative intravenous loading doses of paracetamol on pain after minor hand surgery: Two grams versus one gram. *Acta Chir Belg.* 2010;110(5):529-32. [Link](#)
168. Danelli G, Bonarelli S, Tognu A, et al. Prospective randomized comparison of ultrasound-guided and neurostimulation techniques for continuous interscalene brachial plexus block in patients undergoing coracoacromial ligament repair. *Br J Anaesth.* 2012;108(6):1006-10. [Link](#)
169. Divella M, Cecconi M, Fasano N, et al. Pain relief after total hip replacement: Oral CR oxycodone plus IV paracetamol versus epidural levobupivacaine and sufentanil. A randomized controlled trial. *Minerva Anesthesiol.* 2012;78(5):534-41. [Link](#)
170. Durak P, Yagar S, Uzuner A, Kilic M, Dilber E, Ozgok A. [Postoperative pain therapy after laparoscopic cholecystectomy: Paracetamol versus diclofenac]. *Agri.* 2010;22(3):117-20. [Link](#)
171. Eftekharian H, Tabrizi R, Kazemi H, Nili M. Evaluation of a single dose intravenous paracetamol for pain relief after maxillofacial surgery: A randomized clinical trial study. *J Maxillofac Oral Surg.* 2014;13(4):478-82. [Link](#)
172. Eker HE, Cok OY, Ergenoglu P, Aribogan A, Arslan G. IV paracetamol effect on propofol-ketamine consumption in paediatric patients undergoing ESWL. *J Anesth.* 2012;26(3):351-6. [Link](#)
173. Ekinci M, Ciftci B, Celik EC, Kose EA, Karakaya MA, Ozdenkaya Y. A randomized, placebo-controlled, double-blind study that evaluates efficacy of intravenous ibuprofen and acetaminophen for postoperative pain treatment following laparoscopic cholecystectomy surgery. *J Gastrointest Surg.* 2019;[published online: April 22, 2019] [Link](#)
174. El-Fattah AM, Ramzy E. Pre-emptive triple analgesia protocol for tonsillectomy pain control in children: Double-blind, randomised, controlled, clinical trial. *J Laryngol Otol.* 2013;127(4):383-91. [Link](#)
175. Elseify ZA, El-Khattab SO, Khattab AM, Atta EM, Ajjoub LF. Combined parecoxib and I.V. paracetamol provides additional analgesic effect with better postoperative satisfaction in patients undergoing anterior cruciate ligament reconstruction. *Saudi J Anaesth.* 2011;5(1):45-9. [Link](#)
176. Emir E, Serin S, Erbay RH, Sungurtekin H, Tomatir E. Tramadol versus low dose tramadol-paracetamol for patient controlled analgesia during spinal vertebral surgery. *Kaohsiung J Med Sci.* 2010;26(6):308-15. [Link](#)
177. Erdogan Kayhan G, Sanli M, Ozgul U, Kirteke R, Yologlu S. Comparison of intravenous ibuprofen and acetaminophen for postoperative multimodal pain management in bariatric surgery: A randomized controlled trial. *J Clin Anesth.* 2018;50:5-11. [Link](#)
178. Eremenko AA, Kuslieva EV. [Analgesic and opioid-sparing effects of intravenous paracetamol in the early period after aortocoronary bypass surgery]. *Anesteziol Reanimatol.* 2008;5(5):11-4. [Link](#)

179. Ergenoglu P, Akin S, Yalcin Cok O, et al. Effect of intraoperative paracetamol on catheter-related bladder discomfort: A prospective, randomized, double-blind study. *Curr Ther Res Clin Exp.* 2012;73(6):186-94. [Link](#)
180. Faiz HR, Rahimzadeh P, Visnjevac O, Behzadi B, Ghodratty MR, Nader ND. Intravenous acetaminophen is superior to ketamine for postoperative pain after abdominal hysterectomy: Results of a prospective, randomized, double-blind, multicenter clinical trial. *J Pain Res.* 2014;7:65-70. [Link](#)
181. Faiz SH, Rahimzadeh P, Alebouyeh MR, Sedaghat M. A randomized controlled trial on analgesic effects of intravenous acetaminophen versus dexamethasone after pediatric tonsillectomy. *Iran Red Crescent Med J.* 2013;15(11):e9267. [Link](#)
182. Fenlon S, Collyer J, Giles J, et al. Oral vs intravenous paracetamol for lower third molar extractions under general anaesthesia: Is oral administration inferior? *Br J Anaesth.* 2013;110(3):432-7. [Link](#)
183. Gehling M, Arndt C, Eberhart LH, Koch T, Kruger T, Wulf H. Postoperative analgesia with parecoxib, acetaminophen, and the combination of both: A randomized, double-blind, placebo-controlled trial in patients undergoing thyroid surgery. *Br J Anaesth.* 2010;104(6):761-7. [Link](#)
184. Gokten OE, Kilicarslan H, Dogan HS, Turker G, Kordan Y. Efficacy of levobupivacaine infiltration to nephrostomy tract in combination with intravenous paracetamol on postoperative analgesia in percutaneous nephrolithotomy patients. *J Endourol.* 2011;25(1):35-9. [Link](#)
185. Gousheh SM, Nesioonpour S, Javaher Foroosh F, Akhondzadeh R, Sahafi SA, Alizadeh Z. Intravenous paracetamol for postoperative analgesia in laparoscopic cholecystectomy. *Anesth Pain Med.* 2013;3(1):214-8. [Link](#)
186. Grundmann U, Wornle C, Biedler A, Kreuer S, Wrobel M, Wilhelm W. The efficacy of the non-opioid analgesics parecoxib, paracetamol and metamizol for postoperative pain relief after lumbar microdiscectomy. *Anesth Analg.* 2006;103(1):217-22, table of contents. [Link](#)
187. Gunusen I, Karaman S, Acar A, Sargin A, Firat V. The efficacy of paracetamol versus tenoxicam on postoperative pain and morphine consumption after abdominal hysterectomy: A placebo-controlled, randomized study. *Clin Exp Obstet Gynecol.* 2012;39(1):49-52. [Link](#)
188. Guo H, Wang C, He Y. A meta-analysis evaluates the efficacy of intravenous acetaminophen for pain management in knee or hip arthroplasty. *J Orthop Sci.* 2018; 23(5):793-800. [Link](#)
189. Haddadi S, Marzban S, Karami MS, Heidarzadeh A, Parvizi A, Naderi Nabi B. Comparing the duration of the analgesic effects of intravenous and rectal acetaminophen following tonsillectomy in children. *Anesth Pain Med.* 2014;4(1):e13175. [Link](#)
190. Haliloglu AH, Gokce MI, Tangal S, Boga MS, Tapar H, Aladag E. Comparison of postoperative analgesic efficacy of penile block, caudal block and intravenous paracetamol for circumcision: A prospective randomized study. *Int Braz J Urol.* 2013;39(4):551-7. [Link](#)
191. Ham SY, Song JW, Shim JK, Lee WK, Kim HJ, Kwak YL. Periprocedural analgesic efficacy of a single, pre-emptive administration of propacetamol in catheter ablation for atrial fibrillation: a randomized controlled trial. *Minerva Anesthesiol.* 2018;84(6):675-683. [Link](#)
192. Hamed MA, Al-Saeed MA. The Efficacy of intravenous magnesium sulfate versus intravenous paracetamol on children posttonsillectomy pain and analgesic requirement: a randomized controlled study. *Anesth Essays Res.* 2018;12(3):724-728. [Link](#)
193. Hashemi SM, Esmaeelijah A, Golzari S, et al. Intravenous paracetamol versus patient-controlled analgesia with morphine for the pain management following diagnostic knee arthroscopy in trauma patients: A randomized clinical trial. *Arch Trauma Res.* 2015;4(4):e30788. [Link](#)
194. Hiller A, Helenius I, Nurmi E, et al. Acetaminophen improves analgesia but does not reduce opioid requirement after major spine surgery in children and adolescents. *Spine (Phila Pa 1976).* 2012;37(20):E1225-31. [Link](#)
195. Hong JY, Kim WO, Chung WY, Yun JS, Kil HK. Paracetamol reduces postoperative pain and rescue analgesic demand after robot-assisted endoscopic thyroidectomy by the transaxillary approach. *World J Surg.* 2010;34(3):521-6. [Link](#)
196. Hong JY, Kim WO, Koo BN, Cho JS, Suk EH, Kil HK. Fentanyl-sparing effect of acetaminophen as a mixture of fentanyl in intravenous parent-/nurse-controlled analgesia after pediatric ureteroneocystostomy. *Anesthesiology.* 2010;113(3):672-7. [Link](#)
197. Hong JY, Won Han S, Kim WO, Kil HK. Fentanyl sparing effects of combined ketorolac and acetaminophen for outpatient inguinal hernia repair in children. *J Urol.* 2010;183(4):1551-5. [Link](#)

198. Idehen HO, Edowmonyi NP, Imarengiaye CA, Kute MO. A comparative study of a combination of paracetamol infusion (perfalgan) and intramuscular diclofenac versus intravenous pethidine, in the management of post caesarean pain. *Niger Postgrad Med J.* 2015;22(1):50-5. [Link](#)
199. Jahangiri Fard A, Babaee T, Alavi SM, et al. Intravenous patient-controlled remifentanyl versus paracetamol in post-operative pain management in patients undergoing coronary artery bypass graft surgery. *Anesth Pain Med.* 2014;4(5):e19862. [Link](#)
200. Jokela R, Ahonen J, Seitsonen E, Marjakangas P, Korttila K. The influence of ondansetron on the analgesic effect of acetaminophen after laparoscopic hysterectomy. *Clin Pharmacol Ther.* 2010;87(6):672-8. [Link](#)
201. Juhl GI, Norholt SE, Tonnesen E, Hiesse-Provost O, Jensen TS. Analgesic efficacy and safety of intravenous paracetamol (acetaminophen) administered as a 2 g starting dose following third molar surgery. *Eur J Pain.* 2006;10(4):371-7. [Link](#)
202. Kamali A, Vakilian K, Mortazavi L, Shokrpour M. The effects of Apotel and Remifentanyl on postoperative pain. *Altern Ther Health Med.* [published online January 15, 2018]. [Link](#)
203. Kamel EZ, Abd-Elshafy SK, Sayed JA, Mostafa MM, Seddik MI. Pain alleviation in patients undergoing cardiac surgery; presternal local anesthetic and magnesium infiltration versus conventional intravenous analgesia: a randomized double-blind study. *Korean J Pain.* 2018;31(2):93-101. [Link](#).
204. Kampe S, Warm M, Landwehr S, et al. Clinical equivalence of IV paracetamol compared to IV dipyrone for postoperative analgesia after surgery for breast cancer. *Curr Med Res Opin.* 2006;22(10):1949-54. [Link](#)
205. Kempainen T, Kokki H, Tuomilehto H, Seppa J, Nuutinen J. Acetaminophen is highly effective in pain treatment after endoscopic sinus surgery. *Laryngoscope.* 2006;116(12):2125-8. [Link](#)
206. Khalili G, Salimianfard M, Zarehzadeh A. Comparison between paracetamol, piroxicam, their combination, and placebo in postoperative pain management of upper limb orthopedic surgery (a randomized double blind clinical trial). *Adv Biomed Res.* 2016;5:114. [Link](#).
207. Khalili G, Janghorbani M, Saryazdi H, Emaminejad A. Effect of preemptive and preventive acetaminophen on postoperative pain score: A randomized, double-blind trial of patients undergoing lower extremity surgery. *J Clin Anesth.* 2013;25(3):188-92. [Link](#)
208. Khajavi MR, Sabouri SM, Shariat Moharari R, Pourfakhr P, Najafi A, Etezadi F, Imani F. Multimodal analgesia with ketamine or tramadol in combination with intravenous paracetamol after renal surgery. *Nephrourol Mon.* 2016;8(4):e36491. [Link](#)
209. Kharouba J, Hawash N, Peretz B, Blumer S, Srour Y, Nassar M, Sabbah M, Safadi A, Khorev A, Somri M. Effect of intravenous paracetamol as pre-emptive compared to preventive analgesia in a pediatric dental setting: a prospective randomized study. *Int J Paediatr Dent.* 2018;28(1):83-91. [Link](#)
210. Kilic O, Akand M, Kulaksizoglu H, et al. Intravenous paracetamol for relief of pain during transrectal-ultrasound-guided biopsy of the prostate: A prospective, randomized, double-blind, placebo-controlled study. *Kaohsiung J Med Sci.* 2015;31(11):572-9. [Link](#)
211. Kilicaslan A, Tuncer S, Yuceaktas A, Uyar M, Reisli R. [The effects of intravenous paracetamol on postoperative analgesia and tramadol consumption in cesarean operations]. *Agri.* 2010;22(1):7-12. [Link](#)
212. Kinoshita J, Fushida S, Kaji M, Oyama K, Fujimoto D, Hirono Y, Tsukada T, Fujimura T, Ohyama S, Yabushita K, Kadoya N, Nishijima K, Ohta T. A randomized controlled trial of postoperative intravenous acetaminophen plus thoracic epidural analgesia vs. thoracic epidural analgesia alone after gastrectomy for gastric cancer. *Gastric Cancer.* 2019;22(2):392-402. [Link](#)
213. Ko MJ, Lee JH, Cheong SH, et al. Comparison of the effects of acetaminophen to ketorolac when added to lidocaine for intravenous regional anesthesia. *Korean J Anesthesiol.* 2010;58(4):357-61. [Link](#)
214. Kocum A, Sener M, Izmirli H, Haydardedeoglu B, Aribogan A. [Efficacy of intravenous dexketoprofen trometamol compared to intravenous paracetamol for postoperative pain management after day-case operative hysteroscopy: Randomized, double-blind, placebo-controlled study]. *Agri.* 2014;26(1):15-22. [Link](#)
215. Kocum AI, Sener M, Caliskan E, et al. Intravenous paracetamol and dipyrone for postoperative analgesia after day-case tonsillectomy in children: A prospective, randomized, double blind, placebo controlled study. *Braz J Otorhinolaryngol.* 2013;79(1):89-94. [Link](#)

216. Kokki M, Broms S, Eskelinen M, Neuvonen PJ, Halonen T, Kokki H. The analgesic concentration of oxycodone with co-administration of paracetamol -- a dose-finding study in adult patients undergoing laparoscopic cholecystectomy. *Basic Clin Pharmacol Toxicol.* 2012;111(6):391-5. [Link](#)
217. Kolahdouzan K, Eydi M, Mohammadipour Anvari H, et al. Comparing the efficacy of intravenous acetaminophen and intravenous meperidine in pain relief after outpatient urological surgery. *Anesth Pain Med.* 2014;4(5):e20337. [Link](#)
218. Koppert W, Frotsch K, Huzurudin N, et al. The effects of paracetamol and parecoxib on kidney function in elderly patients undergoing orthopedic surgery. *Anesth Analg.* 2006;103(5):1170-6. [Link](#)
219. Korkmaz Dilmen O, Tunali Y, Cakmakkaya OS, et al. Efficacy of intravenous paracetamol, metamizol and lornoxicam on postoperative pain and morphine consumption after lumbar disc surgery. *Eur J Anaesthesiol.* 2010;27(5):428-32. [Link](#)
220. Koteswara CM, D S. A study on pre-emptive analgesic effect of intravenous paracetamol in functional endoscopic sinus surgeries (FESSs): A randomized, double-blinded clinical study. *J Clin Diagn Res.* 2014;8(1):108-11. [Link](#)
221. Koyuncu O, Hakimoglu S, Ugur M, Akkurt C, Turhanoglu S, Sessler D, Turan A. Acetaminophen reduces acute and persistent incisional pain after hysterectomy. *Ann Ital Chir.* 2018; [published online: May 15, 2018]. [Link](#)
222. Krishna SN, Chauhan S, Bhoi D, Kaushal B, Hasija S, Sangdup T, Bisoi AK. Bilateral erector spinae plane block for acute post-surgical pain in adult cardiac surgical patients: a randomized controlled trial. *J Cardiothorac Vasc Anesth.* 2019; 33(2):368-375. [Link](#)
223. Landwehr S, Kiencke P, Giesecke T, Eggert D, Thumann G, Kampe S. A comparison between IV paracetamol and IV metamizol for postoperative analgesia after retinal surgery. *Curr Med Res Opin.* 2005;21(10):1569-75. [Link](#)
224. Lee SY, Lee WH, Lee EH, Han KC, Ko YK. The effects of paracetamol, ketorolac, and paracetamol plus morphine on pain control after thyroidectomy. *Korean J Pain.* 2010;23(2):124-30. [Link](#)
225. Lee Y, Yu J, Doumouras AG, Ashoorion V, Gmora S, Anvari M, Hong D. Intravenous acetaminophen versus placebo in post-bariatric surgery multimodal pain management: a meta-analysis of randomized controlled trials. *Obes Surg.* 2019;29(4):1420-1428. [Link](#)
226. Machoki MS, Millar AJ, Albetyn H, Cox SG, Thomas J, Numanoglu A. Local anesthetic wound infusion versus standard analgesia in paediatric post-operative pain control. *Pediatr Surg Int.* 2015;31(11):1087-97. [Link](#)
227. Mahajan L, Mittal V, Gupta R, Chhabra H, Vidhan J, Kaur A. Study to compare the effect of oral, rectal, and intravenous infusion of paracetamol for postoperative analgesia in women undergoing cesarean section under spinal anesthesia. *Anesth Essays Res.* 2017;11(3):594-598. [Link](#)
228. Maghsoudi R, Tabatabai M, Radfar MH, et al. Opioid-sparing effect of intravenous paracetamol after percutaneous nephrolithotomy: A double-blind randomized controlled trial. *J Endourol.* 2014;28(1):23-7. [Link](#)
229. Majumdar S, Das A, Kundu R, Mukherjee D, Hazra B, Mitra T. Intravenous paracetamol infusion: Superior pain management and earlier discharge from hospital in patients undergoing palliative head-neck cancer surgery. *Perspect Clin Res.* 2014;5(4):172-7. [Link](#)
230. Marty J, Benhamou D, Chassard D, et al. Effects of single-dose injectable paracetamol versus propacetamol in pain management after minor gynecologic surgery: A multicenter, randomized, double-blind, active-controlled, two-parallel-group study. *Curr Ther Res Clin Exp.* 2005;66(4):294-306. [Link](#)
231. Maund E, McDaid C, Rice S, Wright K, Jenkins B, Woolacott N. Paracetamol and selective and non-selective non-steroidal anti-inflammatory drugs for the reduction in morphine-related side-effects after major surgery: A systematic review. *Br J Anaesth.* 2011;106(3):292-7. [Link](#)
232. Memis D, Inal MT, Kavalci G, Sezer A, Sut N. Intravenous paracetamol reduced the use of opioids, extubation time, and opioid-related adverse effects after major surgery in intensive care unit. *J Crit Care.* 2010;25(3):458-62. [Link](#)
233. Mitra S, Khandelwal P, Sehgal A. Diclofenac-tramadol vs. diclofenac-acetaminophen combinations for pain relief after caesarean section. *Acta Anaesthesiol Scand.* 2012;56(6):706-11. [Link](#)
234. Mohamad AH, McDonnell NJ, Bloor M, Nathan EA, Paech MJ. Parecoxib and paracetamol for pain relief following minor day-stay gynaecological surgery. *Anaesth Intensive Care.* 2014;42(1):43-50. [Link](#)

235. Mollazadeh R, Eftekhari MR, Eslami M. Efficacy of intravenous acetaminophen in periimplantation pain of cardiac electronic devices: a randomized double-blinded study. *J Perianesth Nurs*. 2017;32(3):215-218. [Link](#)
236. Moller PL, Juhl GI, Payen-Champenois C, Skoglund LA. Intravenous acetaminophen (paracetamol): Comparable analgesic efficacy, but better local safety than its prodrug, propacetamol, for postoperative pain after third molar surgery. *Anesth Analg*. 2005;101(1):90-6, table of contents. [Link](#)
237. Moon YE, Lee YK, Lee J, Moon DE. The effects of preoperative intravenous acetaminophen in patients undergoing abdominal hysterectomy. *Arch Gynecol Obstet*. 2011;284(6):1455-60. [Link](#)
238. Mowafi HA, Elmakarim EA, Ismail S, Al-Mahdy M, El-Saflan AE, Elsaid AS. Intravenous lornoxicam is more effective than paracetamol as a supplemental analgesic after lower abdominal surgery: A randomized controlled trial. *World J Surg*. 2012;36(9):2039-44. [Link](#)
239. Murat I, Baujard C, Foussat C, et al. Tolerance and analgesic efficacy of a new i.v. paracetamol solution in children after inguinal hernia repair. *Paediatr Anaesth*. 2005;15(8):663-70. [Link](#)
240. Murata-Ooiwa M, Tsukada S, Wakui M. Intravenous acetaminophen in multimodal pain management for patients undergoing total knee arthroplasty: a randomized, double-blind, placebo-controlled trial. *J Arthroplasty*. 2017;32(10):3024-302. [Link](#)
241. Nikooseresht M, Seifrabiei MA, Davoodi M, Aghajanlou M, Sardari MT. Diclofenac suppository vs. IV acetaminophen combined with IV PCA for postoperative pain management in patients undergoing laminectomy: a randomized, double-blinded clinical trial. *Anesth Pain Med*. 2016;6(3):e36812. [Link](#)
242. Nonaka T, Hara M, Miyamoto C, Sugita M, Yamamoto T. Comparison of the analgesic effect of intravenous acetaminophen with that of flurbiprofen axetil on post-breast surgery pain: A randomized controlled trial. *J Anesth*. 2016;30(3):405-409. [Link](#)
243. Ohnesorge H, Bein B, Hanss R, et al. Paracetamol versus metamizol in the treatment of postoperative pain after breast surgery: A randomized, controlled trial. *Eur J Anaesthesiol*. 2009;26(8):648-53. [Link](#)
244. Olonisakin RP, Amanor-Boadu SD, Akinyemi AO. Morphine-sparing effect of intravenous paracetamol for post operative pain management following gynaecological surgery. *Afr J Med Med Sci*. 2012;41(4):429-36. [Link](#)
245. Oncul AM, Cimen E, Kucukyavuz Z, Cambazoglu M. Postoperative analgesia in orthognathic surgery patients: Diclofenac sodium or paracetamol? *Br J Oral Maxillofac Surg*. 2011;49(2):138-41. [Link](#)
246. Oreskovic Z, Bicanic G, Hrabac P, Tripkovic B, Delimar D. Treatment of postoperative pain after total hip arthroplasty: Comparison between metamizol and paracetamol as adjunctive to opioid analgesics-prospective, double-blind, randomised study. *Arch Orthop Trauma Surg*. 2014;134(5):631-6. [Link](#)
247. Ozmete O, Bali C, Cok OY, Ergenoglu P, Ozyilkan NB, Akin S, Kalayci H, Aribogan A. Preoperative paracetamol improves post-cesarean delivery pain management: a prospective, randomized, double-blind, placebo-controlled trial. *J Clin Anesth*. 2016;33:51-57. [Link](#)
248. Ozkan F, Erdemir F, Erkorkmaz U, Kaya Z, Senayli Y, Parlaktas BS. Comparison of three different analgesic protocols during shockwave lithotripsy. *J Endourol*. 2012;26(6):691-6. [Link](#)
249. Paech MJ, McDonnell NJ, Sinha A, Baber C, Nathan EA. A randomised controlled trial of parecoxib, celecoxib and paracetamol as adjuncts to patient-controlled epidural analgesia after caesarean delivery. *Anaesth Intensive Care*. 2014;42(1):15-22. [Link](#)
250. Parida S, Panga R, Rajappa M, Kundra P. Study of glutathione S-transferase levels in patients receiving intravenous paracetamol perioperatively: a randomized controlled trial. *Indian J Gastroenterol*. 2018;37(6):511-519. [Link](#)
251. Petterson PH, Jakobsson J, Owall A. Intravenous acetaminophen reduced the use of opioids compared with oral administration after coronary artery bypass grafting. *J Cardiothorac Vasc Anesth*. 2005;19(3):306-9. [Link](#)
252. Pickering G, Faure M, Commun F, et al. Tropisetron and paracetamol association in post-operative patients. *Fundam Clin Pharmacol*. 2012;26(3):432-7. [Link](#)
253. Raja S DC, Shetty AP, Subramanian B, Kanna RM, Rajasekaran S. A prospective randomized study to analyze the efficacy of balanced preemptive analgesia in spine surgery. *Spine J*. 2018; [published online: October 22, 2018]. [Link](#)
254. Roofthoof DW, Simons SHP, van Lingen RA, Tibboel D, van den Anker JN, Reiss IKH, van Dijk M. Randomized controlled trial comparing different single doses of intravenous paracetamol for

- placement of peripherally inserted central catheters in preterm infants. *Neonatology*. 2017;112(2):150-158. [Link](#)
255. Rugyte D, Gudaityte J. Intravenous paracetamol in adjunct to intravenous ketoprofen for postoperative pain in children undergoing general surgery: a double-blinded randomized study. *Medicina (Kaunas, Lithuania)*. 2019;55(4):86. [Link](#)
256. Salihoglu Z, Yildirim M, Demiroglu S, et al. Evaluation of intravenous paracetamol administration on postoperative pain and recovery characteristics in patients undergoing laparoscopic cholecystectomy. *Surg Laparosc Endosc Percutan Tech*. 2009;19(4):321-3. [Link](#)
257. Salonen A, Silvola J, Kokki H. Does 1 or 2 g paracetamol added to ketoprofen enhance analgesia in adult tonsillectomy patients? *Acta Anaesthesiol Scand*. 2009;53(9):1200-6. [Link](#)
258. Samulak D, Michalska M, Gaca M, Wilczak M, Mojs E, Chuchracki M. Efficiency of postoperative pain management after gynecologic oncological surgeries with the use of morphine + acetaminophen + ketoprofen versus morphine + metamizol + ketoprofen. *Eur J Gynaecol Oncol*. 2011;32(2):168-70. [Link](#)
259. Sadrolsadat SH, Yousefshahi F, Ostadalipour A, Mohammadi FZ, Makarem J. Effect of intravenous acetaminophen on postoperative pain in vitrectomy: a randomized, double-blind, clinical trial. *Anesth Pain Med*. 2017;7(3):e13639. [Link](#)
260. Sen H, Kulahci Y, Bicerer E, Ozkan S, Dagli G, Turan A. The analgesic effect of paracetamol when added to lidocaine for intravenous regional anesthesia. *Anesth Analg*. 2009;109(4):1327-30. [Link](#)
261. Serclova Z, Dytrych P, Marvan J, et al. Fast-track in open intestinal surgery: Prospective randomized study (clinical trials gov identifier no. NCT00123456). *Clin Nutr*. 2009;28(6):618-24. [Link](#)
262. Shahid M, Manjula BP, Sunil BV. A comparative study of intravenous paracetamol and intravenous tramadol for postoperative analgesia in laparotomies. *Anesth Essays Res*. 2015;9(3):314-319. [Link](#)
263. Shi SB, Wang XB, Song JM, Guo SF, Chen ZX, Wang Y. Efficacy of intravenous acetaminophen in multimodal management for pain relief following total knee arthroplasty: a meta-analysis. *J Orthop Surg Res*. 2018;13(1):250. [Link](#)
264. Shimia M, Parish M, Abedini N. The effect of intravenous paracetamol on postoperative pain after lumbar discectomy. *Asian Spine J*. 2014;8(4):400-4. [Link](#)
265. Silvanto M, Munsterhjelm E, Savolainen S, et al. Effect of 3 g of intravenous paracetamol on postoperative analgesia, platelet function and liver enzymes in patients undergoing tonsillectomy under local anaesthesia. *Acta Anaesthesiol Scand*. 2007;51(9):1147-54. [Link](#)
266. Soltani G, Molkizadeh A, Amini S. Effect of intravenous acetaminophen (paracetamol) on hemodynamic parameters following endotracheal tube intubation and postoperative pain in caesarian section surgeries. *Anesth Pain Med*. 2015;5(6):e30062. [Link](#)
267. Tiippana E, Bachmann M, Kalso E, Pere P. Effect of paracetamol and coxib with or without dexamethasone after laparoscopic cholecystectomy. *Acta Anaesthesiol Scand*. 2008;52(5):673-80. [Link](#)
268. Tunali Y, Akcil EF, Dilmen OK, et al. Efficacy of intravenous paracetamol and dexketoprofen on postoperative pain and morphine consumption after a lumbar disk surgery. *J Neurosurg Anesthesiol*. 2013;25(2):143-7. [Link](#)
269. Tuzuner Oncul AM, Yazicioglu D, Alanoglu Z, Demiralp S, Ozturk A, Ucok C. Postoperative analgesia in impacted third molar surgery: The role of preoperative diclofenac sodium, paracetamol and lornoxicam. *Med Princ Pract*. 2011;20(5):470-6. [Link](#)
270. Unal C, Cakan T, Baltaci B, Basar H. Comparison of analgesic efficacy of intravenous paracetamol and intravenous dexketoprofen trometamol in multimodal analgesia after hysterectomy. *J Res Med Sci*. 2013;18(10):897-903. [Link](#)
271. Unal SS, Aksoy M, Ahiskalioglu A, Erdem AF, Adanur S. The effect of intravenous preemptive paracetamol on postoperative fentanyl consumption in patients undergoing open nephrectomy: A prospective randomized study. *Niger J Clin Pract*. 2015;18(1):68-74. [Link](#)
272. Upadya M, Pushpavathi SH, Seetharam KR. Comparison of intra-peritoneal bupivacaine and intravenous paracetamol for postoperative pain relief after laparoscopic cholecystectomy. *Anesth Essays Res*. 2015;9(1):39-43. [Link](#)
273. Uvarov DN, Orlov MM, Levin AV, Sokolov AV, Nedashkovskii EV. [Role of paracetamol in a balanced postoperative analgesia scheme after thoracotomy]. *Anesteziol Reanimatol*. 2008;4(4):46-9. [Link](#)

274. Uysal HY, Takmaz SA, Yaman F, Baltaci B, Basar H. The efficacy of intravenous paracetamol versus tramadol for postoperative analgesia after adenotonsillectomy in children. *J Clin Anesth.* 2011;23(1):53-7. [Link](#)
275. Uzun S, Erden IA, Canbay O, Aypar U. The effect of intravenous paracetamol for the prevention of rocuronium injection pain. *Kaohsiung J Med Sci.* 2014;30(11):566-9. [Link](#)
276. Uzun S, Aycan IO, Erden IA, Sahin A, Aypar U. The addition of metamizole to morphine and paracetamol improves early postoperative analgesia and patient satisfaction after lumbar disc surgery. *Turk Neurosurg.* 2010;20(3):341-7. [Link](#)
277. Wong I, St John-Green C, Walker SM. Opioid-sparing effects of perioperative paracetamol and nonsteroidal anti-inflammatory drugs (NSAIDs) in children. *Paediatr Anaesth.* 2013;23(6):475-95. [Link](#)
278. Yaghoubi S, Pourfallah R, Barikani A, Kayalha H. The postoperative analgesic effect of morphine and paracetamol in the patients undergoing laparotomy, using PCA method. *Glob J Health Sci.* 2014;6(1):207-14. [Link](#)
279. Yalcin N, Uzun ST, Reisli R, Borazan H, Otelcioglu S. A comparison of ketamine and paracetamol for preventing remifentanyl induced hyperalgesia in patients undergoing total abdominal hysterectomy. *Int J Med Sci.* 2012;9(5):327-33. [Link](#)
280. Yamamoto N, Sakura S, Noda T, Nishiyama A, Dan'ura T, Matsui Y, Ozaki T. Comparison of the postoperative analgesic efficacies of intravenous acetaminophen and fascia iliaca compartment block in hip fracture surgery: A randomised controlled trial. *Injury.* 2019; [published online: March 16, 2019] [Link](#)
281. Yenigun A, Yilmaz S, Dogan R, Goktas SS, Calim M, Ozturan O. Demonstration of analgesic effect of intranasal ketamine and intranasal fentanyl for postoperative pain after pediatric tonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2018;104:182-185. [Link](#)
282. Zeidan A, Mazoit JX, Ali Abdullah M, Maaliki H, Ghattas T, Saifan A. Median effective dose (ED₅₀) of paracetamol and morphine for postoperative pain: A study of interaction. *Br J Anaesth.* 2014;112(1):118-23. [Link](#)
283. Yenigun A, Yilmaz S, Dogan R, Goktas SS, Calim M, Ozturan O. Demonstration of analgesic effect of intranasal ketamine and intranasal fentanyl for postoperative pain after pediatric tonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2018;104:182-185. [Link](#)

US HEOR

284. Agarwal R, Rajanbabu A, Unnikrishnan UG. A retrospective evaluation of the perioperative drug use and comparison of its cost in robotic vs open surgery for endometrial cancer. *J Robot Surg.* 2018; 2(4):665-672 [Link](#).
285. Allen KB, Michael Borkon A, Cohen DJ, Chhatrwalla AK, Aggarwal S, Saxon J, Russell Davis J, Kennedy KF. Intravenous acetaminophen improves outcomes after transapical transcatheter aortic valve replacement. *Innovations (Phila).* 2018;13(4):287-291. [Link](#)
286. Apfel C, Jahr JR, Kelly CL, Ang RY, Oderda GM. Effect of I.V. acetaminophen on total hip or knee replacement surgery: a case-matched evaluation of a national patient database. *Am J Health Syst Pharm.* 2015;72(22):1961-8. [Link](#)
287. Apfel CC, Hornuss C, Souza K, Pergolizzi J, Turan A. Intravenous acetaminophen reduces postoperative nausea and vomiting: a systematic review and meta-analysis [ASA abstract]. *Proc Annu Meet Am Soc Anesthesiol.* 2012.
288. Atkins JR, Titch JF, Norcross WP, Thompson JA, Muckler VC. Preemptive oral acetaminophen for women undergoing total laparoscopic hysterectomy. *Nursing for Women's Health.* 2019;23(2):105-113. [Link](#)
289. Barrington JW, Hansen RN, Lovelace B, Boing EA, Chughtai M, Newman JM, Pham AT, Khlopas A, Sodhi N, Sultan AA, Mont MA. Impact of intravenous acetaminophen on lengths of stay and discharge status after total knee arthroplasty. *J Knee Surg.* 2019;32(1):111-116 [Link](#).
290. Blonk K, Davenport A, Morgan B, Muckler V. Administration of oral acetaminophen to reduce costs for the hysterectomy patient at a community hospital. *J Perianesth Nurs.* 2019;34(1):143-150. [Link](#)

291. Bollinger AJ, Butler PD, Nies MS, Sietsema DL, Jones CB, Endres TJ. Is scheduled intravenous acetaminophen effective in the pain management protocol of geriatric hip fractures?. *Geriatr Orthop Surg Rehabil.* 2015;6(3):202-8. [Link](#)
292. Bourgeois FT, Graham DA, Kesselheim AS, Randolph AG. Cost Implications of escalating intravenous acetaminophen use in children. *JAMA Pediatrics.* 2019; [published online: March 11, 2019] [Link](#)
293. Chidambaran V, Subramanyam R, Ding L, Sadhasivam S, Geisler K, Stubbeman B, Sturm P, Jain V, Eckman MH. Cost-effectiveness of intravenous acetaminophen and ketorolac in adolescents undergoing idiopathic scoliosis surgery. *Paediatr Anaesth.* 2018;28(3):237-248. [Link](#)
294. El Chaar M, Stoltzfus J, Claros L, Wasylik T. IV acetaminophen results in lower hospital costs and emergency room visits following bariatric surgery: a double-blind, prospective, randomized trial in a single accredited bariatric center. *J Gastrointest Surg.* 2016;1(4):715-24. [Link](#)
295. Hansen RN, Pham AT, Boing EA, Lovelace B, Wan GJ, Miller TE. Comparative analysis of length of stay, hospitalization costs, opioid use, and discharge status among spine surgery patients with postoperative pain management including intravenous versus oral acetaminophen. *Curr Med Res Opin.* 2017;33(5):943-948. [Link](#)
296. Hansen RN, Pham A, Strassels SA, Balaban S, Wan GJ. Comparative analysis of length of stay and inpatient costs for orthopedic surgery patients treated with IV acetaminophen and IV opioids vs. IV opioids alone for post-operative pain. *Adv Ther.* 2016;33(9):1635-45. [Link](#)
297. Hansen RN, Pham AT, Boing EA, Lovelace B, Wan GJ, Thomas DA, Fontes ML. Hospitalization costs and resource allocation in cholecystectomy with use of intravenous versus oral acetaminophen. *Curr Med Res Opin.* 2018; 34(9):1549-1555. [Link](#).
298. Hansen RN, Pham AT, Lovelace B, Balaban S, Wan GJ. Comparative analysis of inpatient costs for obstetrics and gynecology surgery patients treated with IV acetaminophen and IV opioids versus IV opioid-only analgesia for postoperative pain. *Ann Pharmacother.* 2017;51(10):834-839. [Link](#)
299. Hansen RN, Pham AT, Boing EA, Lovelace B, Wan GJ, Urman RD. Reduced length of stay and hospitalization costs among inpatient hysterectomy patients with postoperative pain management including IV versus oral acetaminophen. *PLoS One.* 2018;13(9):e0203746. [Link](#)
300. Howell K, Karish C, Bigham M, Metzger K, Thompson M, Christopher J. Saving money for a rainy day: Optimizing intravenous acetaminophen use in the PICU [SCCM abstract 1340]. *Critical Care Medicine.* 2018;46(suppl 1):653.
301. Keller DS, Stulberg JJ, Lawrence JK, Delaney CP. Process control to measure process improvement in colorectal surgery: modifications to an established enhanced recovery pathway. *Dis Colon Rectum.* 2014;57(2):194-200. [Link](#)
302. Kelly JS, Opsha Y, Costello J, Schiller D, Hola ET. Opioid use in knee arthroplasty after receiving intravenous acetaminophen. *Pharmacotherapy.* 2014;34(1):22S26S. [Link](#)
303. Khobrani MA, Camamo JM, Patanwala AE. Effect of Intravenous acetaminophen on post-anesthesia care unit length of stay, opioid consumption, pain, and analgesic drug costs after ambulatory surgery. *P T.* 2017;42(2):125-139. [Link](#)
304. Looke TD, Kluth CT. Effect of preoperative intravenous methocarbamol and intravenous acetaminophen on opioid use after primary total hip and knee replacement. *Orthopedics.* 2013;36(2):25-32. [Link](#)
305. Maiese BA, Pham AT, Shah MV, Eaddy MT, Lunacsek OE, Wan GJ. Hospitalization costs for patients undergoing orthopedic surgery treated with intravenous acetaminophen (IV-APAP) plus other IV analgesics or IV opioid monotherapy for postoperative pain. *Adv Ther.* 2017;34(2):421-435. [Link](#)
306. Mont M, Hansen R, Lovelace B, Khlopas A, Boing E, Barrington J. Intravenous acetaminophen use associated with reduced odds of 30-day readmission after total knee arthroplasty [ASRA abstract

- 4328]. Abstract presented at Annual Pain Medicine Meeting; November 16 – 18, 2017; Lake Buena Vista, FL. 2017.
307. Mont MA, Lovelace B, Pham AT, Hansen RN, Chughtai M, Gwam CU, Khlopas A, Barrington JW. Intravenous acetaminophen may be associated with reduced odds of 30-Day readmission after total knee arthroplasty. *J Knee Surg.* 2019;32(5):414-420. [Link](#)
308. Olbrecht VA, Ding L, Spruance K, Hossain M, Sadhasivam S, Chidambaran V. Intravenous acetaminophen reduces length of stay via mediation of postoperative opioid consumption following posterior spinal fusion in a pediatric cohort. *Clin J Pain.* 2018;34(7):593-599. [Link](#)
309. Pham A, Hansen R, Boing EA, Lovelace B, Wan GJ, Urman R. Reduced length of stay and hospitalization costs among inpatient hysterectomy surgery patients with postoperative pain management including IV versus oral acetaminophen [ASRA abstract 3085]. Abstract Presented at American Society of Regional Anesthesia & Pain Medicine. 2016;November 17-19, 2016:San Diego, CA.
310. Raiff D, Vaughan C, McGee A. Impact of intraoperative acetaminophen administration on postoperative opioid consumption in patients undergoing hip or knee replacement. *Hosp Pharm.* 2014;49(11):1022-32. [Link](#)
311. Shafi S, Collinsworth AW, Copeland LA, Ogola GO, Qiu T, Kouznetsova M, Liao I-C, Mears N, Pham AT, Wan GJ, Masica AL. Association of opioid-related adverse drug events with clinical and cost outcomes among surgical patients in a large integrated health care delivery system. *JAMA Surg.* 2018; 153(8):757-763. [Link](#)
312. Shaffer EE, Woldman RL, Spiegelman A, Strassels SA, Wan GJ, Zimmerman T. Estimating the effect of intravenous acetaminophen (IV-APAP) on length of stay and inpatient costs [ASRA abstract 1364]. Abstract Presented at 41st Annual Regional Anesthesiology & Acute Pain Medicine Meeting. ; March 31 – April 2 2016; New Orleans LA.
313. Shaffer EE, Pham A, Woldman RL, Spiegelman A, Strassels SA Wan GJ, Zimmerman T. Estimating the Effect of intravenous acetaminophen for postoperative pain management on length of stay and inpatient hospital costs. *Adv Ther.* 2017;33(12):2211-2228. [Link](#)
314. Song K, Melroy MJ, Whipple OC. Optimizing multimodal analgesia with intravenous acetaminophen and opioids in postoperative bariatric patients. *Pharmacotherapy.* 2014;34(1):14S21S. [Link](#)
315. Subramanyam R, Varughese A, Kurth CD, Eckman MH. Cost-effectiveness of intravenous acetaminophen for pediatric tonsillectomy. *Paediatr Anaesth.* 2014;24(5):467-475. [Link](#)
316. Urman R, Hansen R, Böing E, Lovelace B, Wan G. Reduced length of stay and hospitalization costs among inpatient appendectomy patients with postoperative pain management including intravenous versus oral acetaminophen [ASRA abstract 4222]. Abstract presented at Annual Pain Medicine Meeting; November 16 – 18, 2017; Lake Buena Vista, FL. 2017.
317. Urman R, Hansen R, Böing E, Lovelace B, Wan G. Reduced length of stay and hospitalization costs among inpatient bariatric patients with postoperative pain management including intravenous versus oral acetaminophen [ASRA abstract 4224]. Abstract presented at Annual Pain Medicine Meeting; November 16 – 18, 2017; Lake Buena Vista, FL. 2017.
318. Urman RD, Boing EA, Khangulov V, Fain R, Nathanson BH, Wan GJ, Lovelace B, Pham AT, Cirillo J. Analysis of predictors of opioid-free analgesia for management of acute post-surgical pain in the United States. *Curr Med Res Opin.* 2019;35(2):283-289. [Link](#)
319. Urman R, Boing E, Pham A, Khangulov V, Fain R, Nathanson B, Zhang X, Wan G, Lovelace B, Cirillo J. Improved outcomes associated with the use of intravenous acetaminophen for management of acute post-surgical pain in cesarean sections and hysterectomies. *J Clin Med Res.* 2018;10(6):499-507. [Link](#)
320. Urman RD, Seger DL, Fiskio JM, Neville BA, Harry EM, Weiner SG, Lovelace B, Fain R, Cirillo J, Schnipper JL. The burden of opioid-related adverse drug events on hospitalized previously opioid-free surgical patients. *J Patient Saf.* 2019; [published online: January 21, 2019]. [Link](#)

321. Wang S, Saha R, Shah N, Hanna A, DeMuro J, Calixte R, Brathwaite C. Effect of intravenous acetaminophen on postoperative opioid use in bariatric surgery patients. *P T*. 2015;40(12):847-850,838. [Link](#)
322. Wasserman I, Poeran J, Zubizarreta N, Babby J, Serban S, Goldberg AT, Greenstein AJ, Memtsoudis SG, Mazumdar M, Leibowitz AB. Impact of intravenous acetaminophen on perioperative opioid utilization and outcomes in open colectomies: a claims database analysis. *Anesthesiology*. 2018; 129(1):77-88. [Link](#).
323. Ziemann-Gimmel P, Hensel P, Koppman J, Marema R. Multimodal analgesia reduces narcotic requirements and antiemetic rescue medication in laparoscopic Roux-en-Y gastric bypass surgery. *Surg Obes Relat Dis*. 2013;9(6):975-80. [Link](#)
324. Zlatev DV, Friedlander DF, Tinay I, Wang Y, Pucheril D, Chung BI, Chang SL. Impact of intravenous acetaminophen on outcomes following radical nephrectomy. [ASCO abstract 663]. *J Clin Oncol*. 2018;36(suppl 6s):663